



## Analysis of Land Training and Aquatic Training on Selected Physical Fitness Variables among Basketball Players

Dr. A. SenthilKumar

Physical Director, Idhaya Engineering College for Women, Chinnasalem, Tamilnadu, India.

Received 19th September 2016, Accepted 20th October 2016

### Abstract

*The purpose of this study was to analyse the aquatic and land training on selected physical fitness variables among Basketball players. To achieve the purpose 30 physically active and interested undergraduate engineering Basketball players were selected as subjects and their age ranged between 18 and 24 years. The subjects are categorized into two groups randomly viz.; Aquatic Training group (ATG), land Training group (LTG) and each group had 15 subjects. The experimental group underwent the experimental treatment for 6 weeks, 3 days per week and a session on each day with 90 minutes duration. Speed, endurance and explosive power were taken as variables for this investigation. Fifty meters run, 12 minutes run and standing vertical jump were tests used to collect the relevant data. The data were collected prior and after the experimental treatment. The collected data was analyzed using analysis of covariance (ANCOVA). The result reveals that aquatic training group showed significant improvement in all the selected physical fitness variables.*

**Keywords:** Aquatic training, Land training, Speed, Endurance, Explosive power, Basketball.

© Copy Right, IJRRAS, 2016. All Rights Reserved.

### Introduction

The benefits of aquatic exercise originate from the supportive nature of the water environment, muscular strengthening and toning of muscles which result from the resistive properties of water as a dense liquid. The buoyancy experienced in water reduces body weight and makes many exercises possible while reducing stress on joints. Buoyancy is the force that water applies in an upward direction against gravity. The buoyant force provided by water decreases the player's weight in relation to the degree of submersion and decreases the amount of force and joint compression during landing. The buoyancy effect of water makes aquatic training an optimal exercise environment for the players and individuals, as impact and stress on joints is reduced (Gappmaier et al., 2006).

Water has several properties that make it an ideal environment for exercise. The buoyancy of water supports the submerged body from the downward pull of gravity, providing up to a 90% reduction in body weight (Darby & Yaekle, 2000; DiPrampo, 1986; Wilder & Brennan, 2004). Benefits of this buoyant effect include less stress and pressure on bone, muscle and connective tissue, while the viscosity and drag force of water provides a resistance proportional to the exerted effort (Wilder & Brennan, 2004). When the velocity of movement doubles, the drag force produced by water

quadruples, providing a resistance training stimulus (Tsourlou, et al., 2006). As the density of water is approximately 800 times that of air (DiPrampo, 1986), the buoyant properties of water reduce forces on the musculoskeletal system, thereby decreasing the risk of overuse injuries such as tendonitis and stress fractures. Aquatic training resulted in similar training effects as land-based training with a possible reduction in stress due to the reduction of impact afforded by the buoyancy and resistance of the water upon landing (Stemm & Bert, 2007). Aquatic exercise does not worsen the joint condition or result in injury (Wang, 2006). The resistance of the water promotes strengthening. Water acts as a variable "accommodating" resistance (Prins, 2009). An aquatic training programme can decrease compression forces, vibration forces and tensional forces that a player may ensure while training on land (Roswell, et al., 2009). The resistance of the water promotes strengthening. In recent years, aquatic training became one of the most important training to improve the physical and physiological variables (Beale et al, 2005). The purpose of this study was to analyse the aquatic and land training on selected physical fitness variables among Basketball players.

### Methods

To achieve these purpose 30 physically active and interested undergraduate Engineering Basketball players were randomly selected as subjects and their age ranged between 18 and 24 years. The subjects are categorized into two groups randomly viz. Aquatic Training group (ATG), Land Training group (CG), and

### Correspondence

Dr.A.Senthilkumar

E-mail: a\_sk7777@yahoo.co.in, Ph. +9176677 12472

each group had 15 subjects. The selected criterion variables Speed was assessed by fifty meter run, Endurance was assessed by coopers'12 minutes run in meters test and explosive power was assessed by Standing Vertical Jump in centimeters. The aquatic training group and land training group underwent the experimental treatment for 6 weeks, 3 days per week and a session on each day with 90 minutes duration.

**Aquatic and land training**

Warming-up exercise was performed in ground and water. After that the land training group and aquatic training group performed the following exercises in the respective land and aquatic places. The water level was just above the hip level. 1. Single leg jump (alternative leg), 2. Double leg jump, 3. High knee action, 4. Aerobic

exercise. These exercises were performed for 90 minutes in a day and for 3 days per week. Initially apilot work was done by the investigators.

**Statistical analysis**

Pre and post test data were collected before and after 6 weeks of training. The collected data was analyzed using analysis of covariance (ANCOVA). The means and standard deviations of both aquatic and land training groups were calculated for speed, endurance and explosive power for the pre as well as post tests. ANCOVA was used to examine significance between testing groups (ATG and LTG). Statistical significance was set to a priority at  $p < 0.05$ . All statistical tests were calculated using the Statistical Package for the Social Science (SPSS) for Windows (Version 15).

**Table 1.** Analysis of covariance for speed of aquatic and land training groups

Test	Aquatic Training Group (ATG)	Land Training Group (LTG)	Sources of Variance	Sum of the Squares	df	Means Squares	F-ratio
Pre test mean	7.46	7.42	BG	0.003	1	0.003	0.039
	0.26	0.27	WG	2.22	28	0.09	
Post test mean	7.08	7.35	BG	0.39	1	0.37	3.75
	0.25	0.38	WG	2.85	28	0.10	
Adjusted post test mean	7.05	7.36	BG	0.46	1	0.47	23.09*
			WG	0.56	27	0.02	


\* Significant at 0.05 level

**Results**

Table I shows that the pre and post test means and standard deviation of aquatic and land training groups on speed. The obtained 'F' value of pre and post test means on speed was 0.039 and 3.75 respectively, which was lesser than table value of 4.19 for degree of freedom 1 and 28 at 0.05 level of confidence; hence there was no significant difference in pre and post test

data of aquatic and land training groups. The analysis of adjusted post test mean data reveals that obtained 'F' value of 23.09 was greater than table of 4.21 for degree of freedom 1 and 27 at 0.05 level of confidence; hence there exist difference in speed among the ATG and LTG groups.

**Figure I.** Diagram showing the pre, post and adjusted post test means on speed



**Table II.** Analysis of covariance for endurance of aquatic and land training groups


Test	Aquatic Training Group (ATG)	Land Training Group (LTG)	Sources of Variance	Sum of the Squares	df	Means Squares	F-ratio
Pre test mean	2239.35	2152.68	BG	56333.35	1	56333.38	0.98
	204.48	274.35	WG	1637186.68	28	58470.96	
Post test mean	2298.68	2165.38	BG	133333.58	1	133333.98	2.35
	209.04	269.99	WG	1558146.86	28	55648.95	
Adjusted post test mean	2256.52	2207.49	BG	17436.26	1	17436.18	50.19*
			WG	9360.12	27	346.68	

\* Significant at 0.05 level

The table II reveals that the pre and post test means and standard deviation of aquatic and land training groups on endurance. The obtained 'F' value of pre and post test means on endurance was 0.98 and 2.35 respectively, which was lesser than table value of 4.19 for degree of freedom 1 and 28 at 0.05 level of confidence; hence there was no significant difference in

pre and post test data of aquatic and land training groups. The analysis of adjusted post test mean data reveals that obtained 'F' value of 50.19 was greater than table value of 4.21 for degree of freedom 1 and 27 at 0.05 level of confidence; hence there exist difference in endurance among the ATG and LTG groups.

**Figure II.** Diagram showing the pre, post and adjusted post test means on Endurance



**Table III.** Analysis of covariance for explosive power of aquatic and land training groups


Test	Aquatic Training Group (ATG)	Land Training Group (LTG)	Sources of Variance	Sum of the Squares	df	Means Squares	F-ratio
Pre test mean	48.02	45.74	BG	38.54	1	38.55	1.28
	4.52	6.32	WG	842.95	28	30.12	
Post test mean	52.16	47.94	BG	128.15	1	128.14	4.55*
	4.59	5.95	WG	789.87	28	28.22	
Adjusted post test mean	50.99	49.00	BG	28.20	1	28.20	24.48*
			WG	31.11	27	1.15	

\* Significant at 0.05 level

The table III indicates that the pre and post test means and standard deviation of aquatic and land training groups on explosive power. The obtained 'F' value of pre test means on explosive power was 1.28 which was lesser than the table value 4.19 for degree of freedom 1 and 28 at 0.05 level of confidence; hence there was no significant difference in pre test

data of aquatic and land training groups. The obtained 'F' value of post test means on explosive power was 4.55, which was greater than the table value 4.19 for degree of freedom 1 and 28 at 0.05 level of confidence; hence there was significant difference in post test data of aquatic and land training groups. The analysis of adjusted post test mean data reveals that obtained 'F' value of 24.48 was greater than table value of 4.21 for degree of freedom 1 and 27 at 0.05 level of confidence; hence there exist difference in explosive power among the ATG and LTG groups.

**Figure III.** diagram showing the pre, post and adjusted post test means on explosive power



### Discussions

The result of the study reveals that there exists significant difference among the groups on all the selected physical fitness variables. The aquatic training group showed significant improvement in all the selected physical fitness variables. Aquatic-based exercise intervention, the resistive properties of water provided a resistive stimulus regardless of whether specific resistance training exercises are prescribed. An aquatic-based exercise demonstrated positive alterations in lower body strength. The majority of these studies suggest that adding deep-water running to an athlete's training regimen has the potential to increase fitness and ultimately improve performance (Burns & Lauder, 2001). The aquatic environment may be used to provide a workload sufficient to create fatigue and produce strength gains in both deconditioned adults and trained athletes (Tsourlou, et.al, 2006). According to Evans and colleagues (1978), the dual effects of buoyancy and resistance make possible high levels of energy expenditure with relatively little movement or strain on lower-joint extremities. Additionally, enhanced temperature regulation during water exercise makes this an ideal environment for obese individuals who have an increased risk of heat intolerance (Wallace, 2003). Martel and co-workers (2005) demonstrated the ability to increase vertical jump in female Basketball players using specific aquatic plyometric training and these improvements could be accomplished with less muscle pain as well.

### Conclusions

Aquatic training group (ATG) showed significant improvement in all selected physical fitness variables namely speed, endurance and explosive power among Basketball players after aquatic based exercise training intervention.

### References

1. Burns, AS and Lauder TD. (2001) Deep water running: an effective non-weight bearing exercise for the maintenance of land-based running performance, *Milit Med* 166: p: 253 – 258.
2. Beale, Angela Krishnan.,(2005).An investigation of the current status of aquatic physical activity in K – 12 public school physical education programmes in the state of Florida; Sport Management, Recreation Management, and Physical Education. Department. 10-04.
3. Darby, L. A., & Yaekle, B. C. (2000). Physiological responses during two types of exercise performed on land and in the water. *Journal of Sports Medicine and Physical Fitness*, 40, 303-311.
4. DiPrampo, P. E. (1986). The energy cost of human locomotion on land and in water. *International Journal of Sports Medicine*, 7, 55-72.
5. Evans, B. W., Cureton, K. J., & Purvis, J. W. (1978). Metabolic and circulatory responses to walking and jogging in water. *Research Quarterly*, 49, 442- 449.
6. Gappmaier, E., Lake, W., Nelson, A. G., & Fisher, A. G. (2006). Aerobic exercise in Water versus

- Walking on Land: Effects on Indices of Fat Reduction and Weight Loss of Obese Women. *Journal of Sports Medicine and Physical Fitness*, 46, 564-569.
7. Martel, G.F., Harmer, M.L., Logan, J.M., and Parker, C.B., (2005). Aquatic plyometric training increases vertical jump in female Basketball players. *Med Sci Sports Exercises*, Vol – 37: p 1814 – 1819.
  8. Prins. Jan H. (2009) Aquatic Rehabilitation, *Serbian Journal of Sport Sciences*.3 (2) Roswell; Greg J. Aaron J. Coutts B; Peter Reaburn; & Stephen Hill-Haas D. (2009). Effects of cold-water immersion on physical performance between successive matches in high-performance junior male soccer players; *Journal of Sports Science* 27(6). 565-573.