



Effect of Aerobic Dance on Selected Physiological and Biochemical Variables of Obese Adolescent Girls

R. Umamaheswari¹ & Dr. A. Mahaboobjan²

¹Ph.D., Research Scholar, Department of Physical Education, Karpagam University, Coimbatore, Tamilnadu, India.

²Professor & Head, Department of Physical Education, Bharathidasan University, Tamilnadu, India.

Received 15th July 2015, Accepted 10th October 2015

Abstract

The purpose of the study was to find out the effect of aerobic dance on selected physiological and biochemical variables among obese adolescent girls. To achieve the purpose of this study fifteen students from Vidya Mandir Institute of Technology, Perundurai, Erode were randomly selected as subjects. They were clinically confirmed cases of obesity and their age ranged between 17 and 19 years. The selected subjects were made to undergo aerobic dance training for the period of 6 weeks. The subjects were free to withdraw their consent in case of feeling any discomfort during the period of their participation, but there was no dropout during the study. The following physiological variables such as resting pulse rate, vital capacity and respiratory rate and biochemical variables such as Low Density Lipoprotein (LDL), High Density Lipoprotein (HDL) and Triglycerides were selected as dependent variables. The selected physiological and biochemical variables were tested with standardized tests. The pre test data were collected before the training programme and the post-test data were collected after the training programme. In both the cases the data were collected in a single day at the same time. Through this study the authors have found out that the selected aerobic dance training improve the selected physiological variables such as resting pulse rate in 8.5%, Respiratory rate in 9.26% and vital capacity in 19.5%, and biochemical variables such as LDL in 2.84%, HDL 19.64% and triglycerides in 4.94%. Hence the authors can recommend the aerobic dance to all the obese adolescent girls for improving the selected physiological and biochemical variables among them.

Keywords: Aerobic Dance, Obese, Adolescent Girls, Physiological, Bio-chemical.

© Copy Right, IJRRAS, 2015. All Rights Reserved.

Introduction

Over the past three decades, the number of young people who are obese has nearly tripled. For the children in the 6-11 age group the statistic for obesity (not overweight) should be 19.6%; for ages 12-19, the statistic for obesity (not overweight) should be 18.1 %;(National Center for Health Statistics, 2010). The consequences of childhood overweight and obesity are serious. Being overweight or obese increases the risk for heart problems, high blood pressure, and other medical problems, and the psychological impact of being overweight can be devastating. Obesity during childhood and adolescence has been associated with higher rates of sickness and death in adulthood, even when adult weight is considered (Must, Jaques, Dallal, Bajema & Dietz, 1992). Aerobic dance is the fitness sport that combines the health and figure benefits of jogging with the fun of dancing. Aerobic dancing is a fun way to get fit. It combines fat burning aerobic movements, muscle building exercises and stretching into routines that are performed according to music. Aerobic dancing is

challenging for college level boys. They feel as though they are performing dance, while firming up their body and strengthening their cardiovascular system. Aerobic dancing is a series of callisthenic exercise movements, accompanied by music. The use of music is a technique of motivation that has been increased in recent years. Aerobic dance is essential to a healthy cardiovascular system. Briefly, aerobic dance is an activity that can be sustained for an extended period of time without building oxygen debt in the muscles. It is a type of dance that overloads the heart and lungs and causes them to work harder than they do when a person is at rest. Aerobic literally means "with air". Aerobic dance is the type of activity in which the amount of oxygen taken in is equal to the amount of oxygen required.

Methodology

The study under investigation was intended to find out the effect of aerobic dance on selected physiological and biochemical variables among obese adolescent girls. It was hypothesized that there might be significant improvement on the selected physiological and biochemical variables among obese adolescent girls due to the effect of aerobic dance training. To achieve the purpose of this study fifteen students from Vidya Mandir

Correspondence

R.Umamaheshwari.

E-mail: umasports@gmail.com, Ph: +9197886 30222

Institute of Technology, Perundurai, Erode were randomly selected as subjects. They were clinically confirmed cases of obesity and their ages ranged between 17 and 19 years. The selected subjects were made to undergo aerobic dance training for the period of 6 weeks. The subjects were free to withdraw their consent in case of feeling any discomfort during the period of their participation, but there was no dropout during the study. The following physiological variables such as resting pulse rate, vital capacity and respiratory rate and biochemical variables such as Low Density Lipoprotein (LDL), High Density Lipoprotein (HDL)

and Triglycerides were selected as dependent variables. The selected physiological and biochemical variables were tested with standardized tests. The pre test data were collected before the training programme and the post-test data were collected after the training programme. In both the cases the data were collected in a single day at the same time.

Results

The collected data were statistically analyzed with dependent ‘t’ test to find out the significant difference between the pre and post tests.


Table I. Table shows the computation of ‘t’ test between pre and post test means of the following variables:

S.No	Variables	Pre test		Post test		't'	Table value
		Mean	SD	Mean	SD		
1	Resting pulse rate	75.33	7.20	68.93	3.26	3.472*	2.09
2	Respiratory rate	16.53	1.25	15.00	1.31	5.002*	2.09
3	Vital capacity	3.00	0.50	4.29	0.68	6.057*	2.09
4	LDL	79.53	9.72	77.27	9.46	4.900*	2.09
5	HDL	51.20	5.05	61.26	9.82	3.729*	2.09
6	Triglycerides	103.80	17.33	98.67	16.12	5.264*	2.09

The above table shows the obtained ‘t’ ratios for pre and post test mean difference in the selected variable of resting pulse rate (3.472), respiratory rate (5.002), vital capacity (6.057), LDL (4.900), HDL (3.729) and triglycerides (5.264). The obtained ratios, when compared with the table value of 2.09 of the degrees of

freedom (1, 19), was found to be statistically significant at 0.05 level of confidence. It was observed that the means gained and losses made from pre to post test were significantly improved the motor fitness and performance variables.

Figure I. Graphical representation showing the mean value of pre and post tests on selected variables



Discussion on Findings

Results of the study shows that the selected physiological variables such as resting pulse rate, respiratory rate and vital capacity and biochemical

variables such as LDL, HDL and triglycerides were improved due to the effect of the aerobic dance training.

Conclusion

Through this study it is found out that the selected aerobic dance training improve the selected physiological variables such as resting pulse rate in 8.5%, respiratory rate in 9.26% and vital capacity in 19.5%, and biochemical variables such as LDL in 2.84%, HDL 19.64% and triglycerides in 4.94%. Hence the aerobic dance can be recommended to all the obese adolescent girls for improving the selected physiological and biochemical variables among them.

References

1. Jan Galen Bishop (2002) "Fitness through Aerobics", Benjamin Cummings, San Francisco.
2. Williams LD, Morton AR. J Sports Sci. (1986) Changes in selected cardio respiratory responses to exercise and in body composition following a 12-week aerobic dance programme. 4(3):189-99.
3. "Statistics Related to Overweight and Obesity". CDC. 2006. Retrieved 2009-01-23.
4. An overview on obesity Emedicine Health. Retrieved on 2010-02-04