



Investigation of the Changes on Selected Speed Related Parameters in Response to Resisted Sprint Training

Dr. N. Aseer Rufus

Assistant Professor, Department of Physical Education, Scott Christian College, Nagercoil, Tamilnadu, India.

Received 15th June 2016, Accepted 2nd August 2016

Abstract

The purpose of the study was to find out the effect of resisted sprint training on speed, stride length and stride frequency among male sprinters. The resisted sprint training exercises included in this training programme was weighted vest, harness running and weighted sled. The response to resisted sprint training was characterized in thirty male sprinters in the age group of eighteen to twenty two years. The selected criterion variables were measured using standard tests and procedures. Analysis of covariance was used to determine the significant difference between pre test and posttest on selected criterion variables. The findings of the study indicate that speed, stride length, stride frequency and anaerobic power increased due to eight weeks of resisted sprint training. These results suggest that the resisted sprint training is an effective means of enhancing the sprinting ability, by bringing out significant changes in speed parameters.

Keywords: Resisted Sprint, Stride Length, Stride Frequency.

© Copy Right, IJRRAS, 2016. All Rights Reserved.

Introduction

Training and technique are very important in developing or improving a sport skill. Generally as the adaptation to training takes place, the efficiency of the skill improves. This can be seen in an improved running style (Martin and Coe, 1991). Speed training, like strength, flexibility and mental skills training have now become an important ingredient in the total programme, particularly where speed of movement is essential in the sport. The aim of resisted sprint training is to condition the athlete to move at high velocity, employing maximal power when needed. In order to do this, the neuromuscular system must be conditioned to very fast movements and training need to be very specific, with a very high anaerobic component. Resisted sprint training can be defined as the systematic addition of weight to the player's body in any form (uniform, vest pants (or) shirts) or to the implements used in sports (sticks, bats, balls and so on). Resisted sprint training is a technique designed to improve explosive concentric movements such as sprinting speed. Resisted sprint training along with strength training speed endurance training, plyometrics and over speed training produce the greatest changes in the exercises fast twitch muscle fibers. Athletes of all ages can use this resisted sprint training programme.

Different resisted speed strategies are in practice including towing, uphill sprints, sand sprints,

and weighted sprints. In fact, resisted towing can involve an athlete towing a weighted sled, tire, speed parachute, or some other device over a set distance (Faccioni, 1994). Various studies have demonstrated that the resisted sprint training can produce significant changes in running speed and running kinematics. However the longer-term training adaptations after resisted sprint training on speed parameters remain unclear. Hence, the investigator was much interested to conduct a study to investigate the effect of resisted sprint training on speed related parameters.

Methodology

Subjects and Variables

For the purpose of this study, thirty inter-collegiate level male sprinters in the age group of 18 to 22 years were recruited, with their consent. The selected dependent variables namely speed, stride length, stride frequency and anaerobic power, were assessed using standard tests and procedures, prior to and immediately after the training protocol. Speed was measured by 50 metres run. While the subjects were allowed to run fast about 50 metres to measure speed the measurement of the length of the stride was taken. The average of two stride of the subject was the individual stride length. Stride frequency was calculated by dividing the number of strides taken by the time elapsed gives the number of strides ran in one-second. The standard instruments such as electronic stopwatches and measuring tape were used.

Correspondence

Dr.N.Aseer Rufus

E-mail: aseerrufus@gmail.com, Ph. +9198651 87847

Training Protocol

The experimental group underwent the resisted sprint training for eight weeks. The resisted sprint training exercises included in this training programme was weighted vest, harness running and weighted sled. More specifically, the training distance comprised of 30-50 meters and the initial intensity was fixed at 75% and it was increased once in two weeks by 5%. The subjects performed these runs with the specified intensity. The rest interval between repetitions was 3 minutes where they stay active and between sets they performed other balance or trunk activities for 10 minutes. The control group did not participate in any specialized training during the period of study.

Experimental Design and Statistical Procedure

The experimental design used for the present investigation was random group design involving thirty subjects. Analysis of covariance (ANCOVA) was used as a statistical technique to determine the significant difference, if any, existing between pretest and posttest data on selected dependent variables. The level of significance was accepted at 0.05 level.

Analysis of the Data

Analysis of covariance was used to determine the impact of resisted sprint training on selected speed related parameters and it is presented in table-I.

Table I. Analysis of Covariance on Selected Speed Parameters of Resisted Sprint Training Group and Control Group


Variables	Groups	Adjusted Mean	SOV	Sum of Squares	df	Mean Square	'F' ratio
Speed	Experimental	7.44	B	2.16	1	2.16	72.00*
	Control	7.86	W	0.75	27	0.03	
Stride Length	Experimental	1.84	B	104.29	1	104.29	30.58*
	Control	1.79	W	92.14	27	3.41	
Stride Frequency	Experimental	3.31	B	0.13	1	0.13	6.50*
	Control	3.20	W	0.46	27	0.02	

Required table value for significance at 0.05 level of confidence for df of 1 and 27 is 4.21

* Significant at 0.05 level.

The findings of the study shows that significant difference existing between resisted sprint training group and control group on speed, stride length, stride frequency and anaerobic power, since the obtained 'F' ratio of 72.00, 30.58, 6.50 and 10.33 respectively were greater than the required table value of 4.21 for significance at 0.05 level of confidence for df of 1 and

27. Hence it is concluded that eight weeks of resisted sprint training can produce significant changes on speed, stride length and stride frequency of male sprinters. The Adjusted Post Test Mean on Selected Speed Parameters of Experimental and Control Group are graphically represented in figure I.



Results and Discussions

Based on the statistical analysis of the data it is concluded that eight weeks of resisted sprint training can produce significant changes on speed, stride length, stride frequency and anaerobic power of male sprinters. Knicker (1997) examined the effects of external resistance on sprinting mechanics and found that even small resistance loads could result in considerable

changes in kinematics and coordination of muscular activity as seen in EMG patterns of lower limb muscles. Zafeiridis and others (2005) found that sprint training with 5 kg sled pulling for 8 weeks improves acceleration performance. It has been well documented by Cavanagh and Kram (1990) that humans increase both stride length and stride frequency as running velocity increases. These results support the previous observation by Eicher (1975)

that, an increase in speed might cause an increase in stride frequency or stride length or either of the parameters. The study undertaken by Medbo and Burgers (1990) reveals that six weeks of training increased the anaerobic capacity by 10%. Hence, it is suggested that competition exercises with additional loads are “the most specific exercises” and important training means. This seems to be that resisted sprint training techniques will somehow impact on the athlete's nervous system to induce greater stride rates. The present data suggest that the greater average running speed in resisted sprint training might be due to increases in stride length.

Conclusions

The result of this study demonstrated that, speed, stride length and stride frequency of male sprinters increased due to eight weeks of resisted sprint training resisted sprint training. Hence it is suggested that the adaptational changes of resisted sprint training exercise are very dynamic and variable to individual.

References

1. Cavanagh P., and Kram, R. (1990). *Biomechanics of Distance Running*, Champaign, IL: Human Kinetics.pp. 35-63.
2. Eicher, Tom. (1975). “Improving Sprinting Speed through Strength Training”, *Athletic Journal*, 65, p .12.
3. Knicker, A., (1997). “Neuromechanics of sprint specific training skills”, In J Wilkerson, K Ludwig & W Zimmermann (Eds.), *Biomechanics in Sports*, XV (pp.17-21). Denton, TX: Texas Woman's University Press.
4. Martin, D., and Coe, P. (1991). *Training Distance Runners* Champaign, IL: Human Kinetics.
5. Medbo, J. I., and Burgers, S. (1990). “Effect of Training on Anaerobic Capacity”, *Journal of Medicine, Science, Sports and Exercise*, 22:4, pp. 501-507.
6. Zafeiridis, A. *et al.*, (2005). “The Effects of Resisted Sled Pulling Sprint Training on Acceleration and Maximum Speed Performance”, *Journal of Sports Medicine Physical Fitness*, 45:3, pp. 284-90.